

INFORIN

Read all of this leaflet carefully before you start taking this medicine.

- **Keep this leaflet. You may need to read it again.**
- **If you have any further questions, ask your doctor or pharmacist.**
- **This medicine has been prescribed for you. Do not pass it on to others.**

It may harm them, even if their symptoms are the same as yours.

INFORIN
Cream 10% (100 mg/1 g)
Ibuprofen

1. WHAT INFORIN IS AND WHAT IT IS USED FOR?

Composition:

1 g cream contains 100 mg ibuprofen.

Excipients: methyl parahydroxybenzoate, propyl parahydroxybenzoate, glycerol, propylene glycol, white soft paraffin, Glyceryl Stearate/Ceteareth-20/ Ceteareth-12/Cetearyl Alcohol/Cetyl Palmitate, paraffin liquid, cetyl alcohol, carbomer, sodium hydroxide, water purified.

Packaging:

Box with tube of 50 g cream (10%).

How INFORIN acts?

Ibuprofen is a derivative of the phenylpropionic acid and belongs to the group of nonsteroidal anti-inflammatory drugs. Applied locally, it acts by reducing the inflammation and pain. The mechanism of the action of ibuprofen is a result of the inhibition of the synthesis and/or the release of prostaglandins (mediators of the inflammation).

When INFORIN is used?

You can use Inforin cream for:

- decreasing of the symptoms of inflammation at rheumatoid arthritis, including juvenile idiopathic arthritis or Still's disease, degenerative rheumatism of the peripheral joints and of the spinal column, other rheumatic diseases of the joints and the soft tissues;
- decreasing of the pain at sporting and other injuries of the locomotory system and soft tissues.

2. WHAT SHOULD YOU KNOW BEFORE YOU USE INFORIN?

When should the medicine not be used?

The medicine should not be used if:

- you are hypersensitive to ibuprofen or some of the excipients of the drug;
- you are hypersensitive to other nonsteroidal anti-inflammatory medicines.

Precautions:

You must be careful with Inforin cream if:

- you suffer from bronchial asthma, and especially if during the using of the medicine appearance of bronchospasm (sudden constriction of the muscles in the walls of bronchioles).

A precaution is necessary at children under 12 years of age.

Do not apply cream on wounds and infected parts of the skin. The cream should not come in contact with mucous membranes of the eyes and other mucous membranes (mouth, nose, genital or anal area).

Use of INFORIN with food and drink:

Does not have influence.

Use during pregnancy and breast-feeding:

Consult with your doctor or pharmacist before you take any medicine.

Inform your doctor if you are pregnant (or you are suspected that you are pregnant) or you are breast-feeding.

Do not use ibuprofen if you are pregnant or breast-feeding.

Effects on ability to drive and use machine:

Does not have influence.

What should you know about excipients that INFORIN contain?

Inforin cream contains preservatives methyl parahydroxybenzoate and propyl parahydroxybenzoate which may cause allergic reactions (possible delayed reaction), propylene glycol which may cause skin irritation, cetyl alcohol which may cause local skin reactions (contact dermatitis).

Taking other medicines:

Please inform your doctor if you are taking, or you will take, any other medicine even those dispensing without prescription.

There are no data for interactions of other drugs and ibuprofen applied locally in the form of cream.

3. HOW TO USE INFORIN?

Your doctor will advise you how long to take the medicine. Do not interrupt the treatment before you are not consult with your doctor.

Thin coat of the cream is applied on the affected area of the body 3-4 times a day with light massage. Should not be applied on the wounds and infected parts of the skin.

If you have impression that effect of INFORIN is too potent or too low, consult with your doctor or pharmacist.

If you take more INFORIN than you should:

There are no data for overdose of ibuprofen applied locally in the form of cream.

If you forget to take INFORIN

If you missed or forgot to apply the medicine, apply the medicine as soon as possible. If the time for the next dose is approach, wait, don't apply the missed dose but the next dose.

Do not apply a double dose to make up for forgotten individual doses!

4. SIDE EFFECTS OF THE MEDICINE:

The local application of ibuprofen may cause local irritation manifested as burning sensation, skin redness and other skin reactions.

If some of the aforementioned side effects appear or you are notice appearance of side effect which is not mentioned, inform your doctor or pharmacist.

5. STORAGE AND SHELF-LIFE:

Storage:

Should be kept at temperature up to 25°C, in original package.

Should be kept out of the reach of children!

Shelf-life:

Printed on the package.

Should not be administered after the expiry date!

Dispensing status:

Over the counter drug (OTC).



Произведено для / Manufactured for:
SPEY MEDICAL Ltd.
7-12 Tavistock Square, London,
WC1H 9LT, England, UK

Производитель / Manufactured by:
REPLEKPHARMA AD.
Kozle st. 188, Skopje,
R Macedonia